

NON-RANDOM PERTURBATIONS OF THE ANDERSON HAMILTONIAN IN THE 1-D CASE.

J. HOLT¹, S. MOLCHANOV², B. VAINBERG³

ABSTRACT. Recently (see [1]), two of the authors applied the Lieb method to the study of the negative spectrum for particular operators of the form $H = H_0 - W$. Here, H_0 is the generator of the positive stochastic (or sub-stochastic) semigroup, $W(x) \geq 0$ and $W(x) \rightarrow 0$ as $x \rightarrow \infty$ on some phase space X . They used the general results in several “exotic” situations, among them the Anderson Hamiltonian H_0 . In the 1-d case, the subject of the present paper, we will prove similar but more precise results.

KEYWORDS: Anderson Hamiltonian, Schrödinger operator, negative eigenvalues

1. INTRODUCTION

The basic Anderson Hamiltonian H_0 has the form $H_0\psi = -\psi'' + V_\omega\psi$ where $V_\omega(x)$ is a nonnegative random potential. We'll consider here V_ω in the following way. Let $x_n \in \mathbb{R}$ be an increasing sequence and consider a partition of $\mathbb{R}$ into intervals I_k defined by $I_k = \{x : -l \leq x - x_k \leq l\}$ for $k \in \mathbb{Z}$ and positive l . Let $L_k = x_{k+1} - x_k - 2l$ represent the distance between consecutive intervals I_k . We'll suppose that $L_k(\omega) \geq 0, \omega \in (\Omega, \mathcal{F}, P)$ are independent and identically distributed random variables which are unbounded from above, with some density $p(x)$ and finite expectation

$$\mathbb{E}[L_k] = \int_0^\infty xp(x) dx.$$

Define the random potential V_ω by

$$V_\omega(x) = \sum_{k \in \mathbb{Z}} h\mathbb{I}_k(x)$$

¹ University of South Carolina Lancaster, Lancaster, SC 29721, USA. email: jholt@mailbox.sc.edu.

² University of North Carolina Charlotte, Charlotte, NC 28223, USA. email: smolchan@uncc.edu.

³ University of North Carolina Charlotte, Charlotte, NC 28223, USA. email: brvainbe@uncc.edu.

where $\mathbb{I}_k$ is the indicator function of the interval I_k and $h > 0$. It is easy to see that $H_0 \geq 0$ and that each realization P -a.s contains arbitrarily long intervals where $V_\omega = 0$. This implies that $\text{Sp}(H_0) = [0, \infty)$ P -a.s. Let $W(x)$ be a continuous function on $\mathbb{R}$ such that $W(x) \geq 0$ and $W(x) \rightarrow 0$ as $|x| \rightarrow \infty$. Then the negative spectrum $\text{Sp}(H)$ of the non-random perturbation of H_0 defined by $H = H_0 - W$ is discrete with a possible accumulation point only at 0, and the random variable $\mathcal{N}_0(W) = \#\{\lambda_i(\omega) < 0\}$ can be either finite or infinite. Here, $\{\lambda_i\} = \text{Sp}(H) \cap (-\infty, 0)$ so that $\mathcal{N}_0(W)$ represents the number of negative eigenvalues. Note that the event $\{\omega \in \Omega : \mathcal{N}_0(W) = \infty\}$ belongs to the tail σ -algebra of the potential V_ω and therefore the Kolmogorov zero-one law implies that either $P\{\mathcal{N}_0(W) < \infty\} = 1$, or $P\{\mathcal{N}_0(W) < \infty\} = 0$. Our goal in this work is to find conditions on W for which $P\{\mathcal{N}_0 < \infty\} = 1$ and $\mathbb{E}[\mathcal{N}_0] < \infty$. In particular, we'll establish the following fact for a Bernoulli potential as a consequence of our general results on the wider class of Kronig-Penney type potentials. Such results are related to the classical renewal process in reliability theory.

Theorem 1.1. *For $h > 0$ and any $0 < p < 1$ let $V_\omega = h \sum_{k \in \mathbb{Z}} \epsilon_k(\omega) \mathcal{X}_k$ where $\{\epsilon_k(\omega)\}$ are independent and identically distributed with $P(\epsilon_k = 1) = p$ and $P(\epsilon_k = 0) = 1 - p$. Here $\mathcal{X}_k$ is the indicator of the interval $[k - 1/2, k + 1/2)$. Then if $W(x) < C_p / \ln^2(|x| + 1)$ for large enough x , then $P\{\mathcal{N}_0(W) < \infty\} = 1$, while $P\{\mathcal{N}_0(W) < \infty\} = 0$ if $W(x) > C_p / \ln^2(|x| + 1)$. Here, C_p is the constant $C_p = \log_{1/p}^2 \pi^2$.*

Our results here are similar to the d -dimensional results obtained in [1]. However, whereas we were able to establish a distinct borderline between finitely many and infinitely many negative eigenvalues in the one dimensional case by finding C_p exactly, the result in [1] only shows the following:

Theorem 1.2. *Let $V_\omega = h \sum_{k \in \mathbb{Z}^d} \epsilon_k(\omega) \mathcal{X}_k$ where $\mathcal{X}_k$ is the indicator of the cube $\{(x_1, \dots, x_d) : -1/2 \leq x_i - k_i < 1/2\}$. There exist positive constants $C_- < C_+$ so that if $0 \leq W(x) \leq C_- / \ln^2(|x| + 2)$ then $P\{\mathcal{N}_0(W) < \infty\} = 1$ while if $W(x) \geq C_+ / \ln^2(|x| + 2)$ then $P\{\mathcal{N}_0(W) = \infty\} = 1$.*

Remark. Estimations of the constants C_- and C_+ in the above theorem are not known as they are related to difficult problems in percolation theory.

Remark. Let's note that $V_\omega(x)$ can be viewed as an asymptotically stationary “renewal” process in which L_k can be thought of as the time between successive occurrences of the event $V_\omega(x) = h$.

Additionally we formulate and prove two more results showing a clear relationship between the tails $P(L_k > x)$ and the function W , and which also demonstrate a distinct borderline between finite and infinitely many eigenvalues. The first of these results when L_k have very “light tails”, that is $P(L_k > x) \sim \exp(-\eta x^\alpha/\alpha)$ as $x \rightarrow \infty$ where $\eta, \alpha > 0$, is actually a generalization of Theorem 1.1. In fact, Theorem 1.1 can be obtained by considering the case $\alpha = 1$.

Theorem 1.3. *Suppose $P\{L_k > x\} \sim e^{-\eta x^\alpha/\alpha}$ as $x \rightarrow \infty$ for some $\eta, \alpha > 0$. Let $C_{\eta,\alpha} = (\eta\alpha^{-1})^{2/\alpha}\pi^2$. If for large k $w_k < C_{\eta,\alpha}/\ln^{2/\alpha} k$, then $\mathcal{N}_0(W) < \infty$ P -a.s. On the other hand, if for large k , $w_k > C_{\eta,\alpha}/\ln^{2/\alpha} k$, then $\mathcal{N}_0(W) = \infty$ P -a.s.*

Remark. It shows that the borderline between P -a.s finiteness of $\mathcal{N}_0(W)$ goes along the line

$$w_k \sim \frac{C_{\eta,\alpha}}{\ln^{2/\alpha} k}.$$

Theorem 1.4. *Suppose $P(L_k > x) \sim c/x^\alpha$ as $x \rightarrow \infty$ with $\alpha > 1$. If for large k , $w_k < t^{-\beta}$ with $\beta > 2/\alpha$, then $\mathcal{N}_0(W) < \infty$ P -a.s. and $\mathbb{E}[\mathcal{N}_0(W)] < \infty$. If on the other hand, $\beta < 2/\alpha$, then $\mathcal{N}_0(W) = \infty$ P -a.s. and $\mathbb{E}[\mathcal{N}_0(W)] = \infty$. It shows that the borderline between P -a.s finiteness of $\mathcal{N}_0(W)$ and the expectation $\mathbb{E}[\mathcal{N}_0(W)]$ goes along the line*

$$w_k \sim \frac{1}{k^{2/\alpha}}.$$

Remark. Both of these asymptotic equivalences are corollaries of more general results established in Theorems 4.1 and 4.2.

It is helpful to first consider the Anderson Hamiltonian H_0 on the half axis defined by

$$H_0\psi = -\psi'' + V_\omega\psi, \quad \psi(0) = 0$$

and the case where $h = \infty$ and W constant on the intervals $I_k = [x_k + l, x_{k+1} - l]$. Let $W(x) = w_k$ for $x \in I_k$. With $h = \infty$, we must interpret H as a direct sum of operators $H_k = -d^2/dx^2 - w_k\mathbb{I}_k$ on a system of decoupled wells $[x_k + l, x_{k+1} - l]$ with Dirichlet boundary conditions at the edges. For each k , the spectrum of H_k is discrete

with eigenvalues

$$\lambda_{k,i} = \frac{\pi^2 i^2}{L_k^2} - w_k$$

and $\mathcal{N}_{0,k}(W) = \#\{\lambda_{k,i} < 0\} = \left\lfloor \frac{\sqrt{w_k} L_k}{\pi} \right\rfloor$. The total number of negative eigenvalues for H is then

$$\mathcal{N}_0(W) = \sum_{k=1}^{\infty} \left\lfloor \frac{\sqrt{w_k} L_k}{\pi} \right\rfloor.$$

This series of independent random variables is finite if and only if the series contains finitely many non-zero terms. But,

$$P \left\{ \left\lfloor \frac{\sqrt{w_k} L_k}{\pi} \right\rfloor \geq 1 \right\} = P \left\{ L_k > \frac{\pi}{\sqrt{w_k}} \right\} = \int_{\pi/\sqrt{w_k}}^{\infty} p(x) dx.$$

Due to the Borel-Cantelli lemmas, we have the following proposition:

Proposition 1.5. $\mathcal{N}_0(W) < \infty$ *P*-a.s. if and only if

$$\sum_{k=1}^{\infty} \int_{\pi/\sqrt{w_k}}^{\infty} p(x) dx = \sum_{k=1}^{\infty} P(L_1 > \pi/\sqrt{w_k}) < \infty.$$

Considering the expectation for $\mathcal{N}_0(W)$, it is clear that

$$\int_{\pi/\sqrt{w_k}}^{\infty} \left(\frac{\sqrt{w_k}}{\pi} x - 1 \right) p(x) dx \leq \mathbb{E}[\mathcal{N}_{0,k}] \leq \int_{\pi/\sqrt{w_k}}^{\infty} \frac{\sqrt{w_k}}{\pi} x p(x) dx \quad (1)$$

since

$$\mathbb{E}[\mathcal{N}_{0,k}] = \sum_{j \geq 1} P(\mathcal{N}_{0,k} \geq j) = \sum_{j \geq 1} P \left(L_k > \frac{j\sqrt{w_k}}{\pi} \right).$$

With $F_k(x) = P(L_k > x)$ and the condition of finiteness of the expectation of L_k , it follows by integration by parts that

$$\int_{\pi/\sqrt{w_k}}^{\infty} x p(x) dx = \int_{\pi/\sqrt{w_k}}^{\infty} F_k(x) dx + \frac{\pi}{\sqrt{w_k}} F_k \left(\frac{\pi}{\sqrt{w_k}} \right)$$

so that

$$\frac{\sqrt{w_k}}{\pi} \int_{\pi/\sqrt{w_k}}^{\infty} F_k(x) dx \leq \mathbb{E}[\mathcal{N}_{0,k}] \leq \frac{\sqrt{w_k}}{\pi} \int_{\pi/\sqrt{w_k}}^{\infty} F_k(x) dx + F_k \left(\frac{\pi}{\sqrt{w_k}} \right).$$

It gives immediately necessary and sufficient conditions for the finiteness of the expectation.

Before considering the general case, we present a few examples to demonstrate how the borderline behavior between finitely many and

infinitely many negative eigenvalues depend on the decay of W and the bump separation lengths L_k .

Example 1.6. *Let's consider the situation when $L_k \geq 0$ are distributed with exponential tails, that is $P\{L_k > x\} = e^{-\eta x}$ for some $\eta > 0$. If for large k $w_k < \frac{\eta^2 \pi^2}{\ln^2 k}$, then $\mathcal{N}_0(W) < \infty$ P -a.s. On the other hand, if for large k , $w_k > \frac{\eta^2 \pi^2}{\ln^2 k}$, then $\mathcal{N}_0(W) = \infty$ P -a.s. It shows that the borderline between P -a.s finiteness of $\mathcal{N}_0(W)$ goes along the line*

$$w_k \sim \frac{\eta^2 \pi^2}{\ln^2 k}.$$

Remark. Actually, we can say even more in the previous example. Again, assuming that $P\{L_k \geq x\} = e^{-\eta x}$, $\mathbb{E}[L_k] = 1$ and $P(L_k \geq \pi w_k^{-1/2}) = \exp(-\eta \pi w_k^{-1/2})$. These exponential tails are given by the density function $p(x) = \eta e^{-\eta x}$. Therefore by (1)

$$\frac{\sqrt{w_k}}{\eta \pi} e^{-\eta \pi / \sqrt{w_k}} \leq \mathbb{E}[\mathcal{N}_{0,k}] \leq \left(\frac{\sqrt{w_k}}{\eta \pi} + 1 \right) e^{-\eta \pi / \sqrt{w_k}} \quad (2)$$

as $w_k \rightarrow 0$. So, for small $\delta > 0$ and

$$w_k = \frac{\eta^2 \pi^2 (1 - \delta)^2}{\ln^2 k}$$

it follows that $\mathbb{E}[\mathcal{N}_{0,k}] = O(n^{-1-\beta})$ where $1 + \beta = (1 - \delta)^{-1}$. Consequently $\mathcal{N}_0(W) = \sum_k \mathbb{E}[\mathcal{N}_{0,k}] < \infty$. If instead

$$w_k = \frac{\eta^2 \pi^2 (1 + \delta)^2}{\ln^2 k}$$

then by (2)

$$\mathbb{E}[\mathcal{N}_{0,k}] \geq \frac{C_\delta}{k^{1-\beta} \ln k}$$

with $1 - \beta = (1 + \delta)^{-1}$ and consequently $\mathbb{E}[\mathcal{N}_0(W)] = \infty$. This shows that the decay in W and the borderline between $\mathbb{E}[\mathcal{N}_0(W)] < \infty$ and $\mathbb{E}[\mathcal{N}_0(W)] = \infty$ is also along

$$W(x) \sim \frac{\eta^2 \pi^2}{\ln^2 x}.$$

Example 1.7. *Let's consider the heavy tailed case where $P(L_k > x) \sim c/x^\alpha$ as $x \rightarrow \infty$ with $\alpha > 2$. If for large k , $w_k < t^{-\beta}$ with $\beta > 2/\alpha$, then $\mathcal{N}_0(W) < \infty$ P -a.s. and $\mathbb{E}[\mathcal{N}_0(W)] < \infty$. If on the other hand,*

$\beta < 2/\alpha$, then $\mathcal{N}_0(W) = \infty$ *P*-a.s. and $\mathbb{E}[\mathcal{N}_0(W)] = \infty$. It shows that the borderline between *P*-a.s finiteness of $\mathcal{N}_0(W)$ and the expectation $\mathbb{E}[\mathcal{N}_0(W)]$ goes along the line

$$w_k \sim \frac{1}{k^{2/(\alpha-1)}}.$$

2. DESCRIPTION OF THE MODEL IN THE GENERAL CASE

One can split the real line $\mathbb{R}$ into two half axes, and introduce a Dirichlet boundary condition at the point of splitting, and reduce the initial spectral problem on $\mathbb{R}$ to the union of two similar problems on the half axes. Consider the 1-D Schrödinger operator $H_0\psi = -\psi'' + V_\omega(x)\psi$ with boundary condition $\psi(0) = 0$ for $x \geq 0$. The random potential V_ω is defined by

$$V_\omega(x) = \sum_{k \in \mathbb{Z}} h \mathbb{I}_k(x, \omega)$$

where $h > 0$, $\mathbb{I}_k$ is the indicator of the interval $I_k = [x_k - l, x_k + l]$ and $x_k > 0$ is a random increasing sequence of real numbers. A typical realization of V_ω is given in Figure 1 below, and consists of bumps of constant height $h > 0$ and width $2l > 0$, separated by wells of random length $L_k = x_{k+1} - x_k - 2l$ where the potential $V_\omega = 0$. We will assume that L_k are independent and identically distributed with some density $p(x)$ and finite expectation

$$\mathbb{E}[L_k] = \int_0^\infty xp(x) dx < \infty$$

and that for all $t > 0$, $P(L_k > t) > 0$. Let's note that the condition $\mathbb{E}[L_k] < \infty$ implies that $V(x, \omega)$ is stationary near the point ∞ .




Figure 1.

3. DIRICHLET-NEUMANN BRACKETING

Consider the partition of $\mathbb{R}^+$ defined by the set $P = \{0, x_1, x_2, \dots\}$. Let H_D be the operator on $L^2(\mathbb{R}^+)$ defined by $H_D\psi = -\psi'' + V_\omega\psi$ with Dirichlet boundary conditions at each point of P , that is, $\psi(x_k) = 0$. Similarly, define the operator H_N by $H_N\psi = -\psi'' + V_\omega\psi$ with Neumann boundary conditions at each point of P , that is, $\psi'(x_k) = 0$. Then, both H_D and H_N are direct sums of operators on finite intervals $I_k = [x_k, x_{k+1}]$. Denote by $\mathcal{N}_{0,D}(W)$, the number of negative eigenvalues for the operator $\mathcal{H}_D \equiv H_D - W$, and by $\mathcal{N}_{0,N}(W)$, the number of negative eigenvalues for the operator $\mathcal{H}_N \equiv H_N - W$ (we will also occasionally use the notations $\mathcal{N}_0(\mathcal{H}_D)$ and $\mathcal{N}_0(\mathcal{H}_N)$ for the numbers of negative eigenvalues of $\mathcal{H}_D$ and $\mathcal{H}_N$). The following classical fact due to H. Weyl (so called Dirichlet-Neumann bracketing, see [2]) will be central in our calculations.

Lemma 3.1. *Let $H = H_0 - W$ and $\mathcal{N}_0(W) = \#\{\lambda_i < 0\}$. Then $\mathcal{N}_{0,D}(W) \leq \mathcal{N}_0(W) \leq \mathcal{N}_{0,N}(W)$.*

Let's take a closer look at the operators $\mathcal{H}_D$ and $\mathcal{H}_N$. Since H_D and H_N are direct sums of operators on the intervals I_k , so are $\mathcal{H}_D$ and $\mathcal{H}_N$. Denote these operators respectively by $\mathcal{H}_{D,k}$ and $\mathcal{H}_{N,k}$. Specifically, $\mathcal{H}_{D,k}$ and $\mathcal{H}_{N,k}$ are defined by the same differential expression as $\mathcal{H}_D$ and $\mathcal{H}_N$, but with Dirichlet boundary conditions at the edges of I_k in the case of $\mathcal{H}_{D,k}$ and Neumann boundary conditions in the case of $\mathcal{H}_{N,k}$. In this case, the numbers of negative eigenvalues for $\mathcal{H}_D$ and $\mathcal{H}_N$ are the sum total of the numbers of negative eigenvalues for the operators $\mathcal{H}_{D,k}$ and $\mathcal{H}_{N,k}$, that is,

$$\mathcal{N}_0(\mathcal{H}_D) = \sum_k \mathcal{N}_0(\mathcal{H}_{D,k}), \quad \mathcal{N}_0(\mathcal{H}_N) = \sum_k \mathcal{N}_0(\mathcal{H}_{N,k}).$$

Now, the spectrum of $\mathcal{H}_{D,k}$ and $\mathcal{H}_{N,k}$ are discrete for all k , with random eigenvalues $\{\lambda_{n,k}^{(D)} : n \geq 0\}$ and $\{\lambda_{n,k}^{(N)} : n \geq 0\}$ accumulating only to ∞ .

The following definition will be important in what follows. Two functions $W^+(x)$ and $W^-(x)$ will be called upper and lower approximations for $W(x)$ if for some $z_0 > 0$, $W^-(x) \leq W(x) \leq W^+(x)$ whenever $x \geq z_0$. We will construct them in the following way. Let us note that under the assumptions on L_i we have $x_k = L_1 + \dots + L_k + (2k-1)l$ for each k and by the strong law of large numbers, $x_k/k \rightarrow \alpha$ P -a.s. where $\alpha = \mathbb{E}[L_i] + 2l$. Therefore, given $\varepsilon > 0$ there exists $k_0 = k_0(\varepsilon, \omega)$ such that when $k \geq k_0$, $(1 - \varepsilon)\alpha k \leq x_k \leq (1 + \varepsilon)\alpha k$. Now, for any

$x \geq x_{k_0}$ there is some k for which $x \in I_k$, and since W is monotonically decreasing,

$$W((1 + \varepsilon)\alpha(k + 1)) \leq W(x_{k+1}) \leq W(x) \leq W(x_k) \leq W((1 - \varepsilon)\alpha k).$$

So, for any $\varepsilon > 0$, the functions W_ε^+ and W_ε^- defined as $W_\varepsilon^+(x) \equiv \sum_{k=1}^\infty W((1 - \varepsilon)\alpha k)\mathbb{I}_k(x)$ and $W_\varepsilon^-(x) \equiv \sum_{k=1}^\infty W((1 + \varepsilon)\alpha(k + 1))\mathbb{I}_k(x)$ are piecewise constant functions for which $W_\varepsilon^-(x) \leq W(x) \leq W_\varepsilon^+(x)$ whenever $x \geq x_{k_0}$. Thus, W_ε^+ and W_ε^- are upper and lower approximations for W .

By the Sturm comparison and oscillation theorems (see [3]) it immediately follows that

$$\mathcal{N}_{0,D,k}(W_\varepsilon^-) \leq \mathcal{N}_{0,D,k}(W) \leq \mathcal{N}_{0,D,k}(W_\varepsilon^+) \quad (3)$$

when $k \geq k_0(\varepsilon, \omega)$. Further, since W_ε^- , W_ε^+ are constant on intervals I_k , the spectra $\sigma(H_{\cdot,k} - W_\varepsilon^-(x))$ and $\sigma(H_{\cdot,k} - W_\varepsilon^+(x))$ can be determined exactly (see below). We first consider a simple example, the results of which will be used in the next section.

Example 3.2. *Let's consider the operators $h_N\psi = -\psi'' + V(x)\psi$ and $h_D\psi = -\psi'' + V(x)\psi$ with Dirichlet boundary conditions at $-L-l$ and $L+l$ in the case of h_D and Neumann conditions in the case of h_N . Let V be the symmetric function defined by $V = h(\mathbb{I}_{[-L-l, -L]} + \mathbb{I}_{[L, L+l]})$:*

Both h_N and h_D have discrete spectrum bounded below by 0, with corresponding eigenfunctions $\psi_i^{(D)}$ and $\psi_i^{(N)}$ for $i \geq 0$. For even i , these eigenfunctions are even, while odd for odd i . This is due to the fact that V is symmetric. We begin by estimating the lowest eigenvalue $\mu_0^{(D)}$ of h_D . Of course, the lowest eigenvalue corresponds to the eigenfunction $\psi_0^{(D)}$. Set $\mu^{(D)} = k^2$. On the interval $[-(L+l), -L]$, the eigenfunction $\psi^{(D)}(x)$ is given by

$$\psi^{(D)}(x) = A \sinh \sqrt{h - k^2}(x + L + l)$$

while on the interval $[-L, 0]$,

$$\psi^{(D)}(x) = B \cos(kx).$$

The continuity conditions, $\psi^{(D)}(-L-0) = \psi^{(D)}(-L+0)$ and $\psi'(-L-0) = \psi'(-L+0)$ immediately provide the eigenvalue equation,

$$k \tan(kL) = \sqrt{h - k^2} \coth(\sqrt{h - k^2}l). \quad (4)$$

Looking ahead to the “real” model, we solve the above equation for $L \rightarrow \infty$. Equation (4) then implies $k_0 \rightarrow 0$ and that for some $\delta_0 > 0$

$$k_0 L = \frac{\pi}{2} - \delta_0 \quad (5)$$

The right side of (4) is analytic in k , and after expansion as a Taylor series, we have

$$\tan(kL) = \frac{A_{0,D}}{k} + A_{1,D}k + O(k^3)$$

or equivalently

$$\cot(kL) = B_{0,D}k - B_{1,D}k^3 + O(k^5) \quad (6)$$

where $A_{j,D}$ and $B_{j,D}$ are constants depending only on h and l . In particular, $B_{0,D} = 1/A_{0,D}$ and $B_{1,D} = A_{1,D}/A_{0,D}$ where $A_{0,D} = \sqrt{h} \coth(\sqrt{h}l)$ and $A_{1,D} > 0$. Substitution of (5) into (6) gives

$$\tan(\delta_0) = B_{0,D}k_0 - B_{1,D}k_0^3 + O(k_0^5)$$

which for $\delta_0 \rightarrow 0$ implies

$$\delta_0 = B_{0,D}k_0 - B_{1,D}k_0^3 + O(k_0^4)$$

or in terms of k_0

$$k_0 = \frac{\pi}{2L} - \frac{1}{L} (B_{0,D}k_0 - B_{1,D}k_0^3) + O(k_0^4). \quad (7)$$

Iterating (7) in k_0 gives

$$\sqrt{\mu_0^{(D)}} = \frac{\pi}{2L} \left(1 - \frac{B_{0,D}}{L} \right) + O\left(\frac{1}{L^3}\right) \quad (8)$$

Similar steps give also the lowest eigenvalue for h_N . In fact, the eigenvalue equation for h_N is

$$k \tan(kL) = \sqrt{h - k^2} \tanh(\sqrt{h - k^2}l) \quad (9)$$

and gives the lowest eigenvalue

$$\sqrt{\mu_0^{(N)}} = \frac{\pi}{2L} \left(1 - \frac{B_{0,N}}{L} \right) + O\left(\frac{1}{L^3}\right) \quad (10)$$

where $B_{0,N}$ is a positive constant depending only on h and l .

Remark. It is interesting to note that the lowest eigenvalue for h_N is asymptotically equivalent to the lowest eigenvalue for h_D .

4. THE CENTRAL THEOREMS

The following results will be important in the study of the borderline decay of $W(x)$.

Given $\varepsilon > 0$, set $w_{k,\varepsilon}^+ = W((1 - \varepsilon)\alpha k)$, $w_{k,\varepsilon}^- = W((1 + \varepsilon)\alpha k)$, and define the upper and lower approximations $W_\varepsilon^+(x) = \sum_{k=1}^\infty w_{k,\varepsilon}^+ \mathbb{I}_k(x)$ and $W_\varepsilon^-(x) = \sum_{k=1}^\infty w_{k,\varepsilon}^- \mathbb{I}_k(x)$. Where $\mathbb{I}_k$ are the indicator functions of the intervals $I_k = [x_k, x_{k+1}]$. Define the operators $H_{D,k} = -d^2/dx^2 + V_k(x)$ with Dirichlet boundary conditions at x_k and x_{k+1} and $H_{N,k} = -d^2/dx^2 + V_k(x)$ with Neumann boundary conditions at x_k and x_{k+1} . Here $V_k = \mathbb{I}_{[x_k, x_{k+1}]} + \mathbb{I}_{[x_{k+1}-l, x_{k+1}]}$. Set $\mathcal{H}_{D,k,\varepsilon} = H_{D,k} - W_\varepsilon^- \mathbb{I}_k$ and $\mathcal{H}_{N,k,\varepsilon} = H_{N,k} - W_\varepsilon^+ \mathbb{I}_k$. Finally $\mu_{0,k}^{(D)}$ and $\mu_{0,k}^{(N)}$ denote respectively, the lowest eigenvalues of $H_{D,k}$ and $H_{N,k}$.

Theorem 4.1. *Let $\varepsilon > 0$ be given. If*

$$\sum_{k=1}^\infty P\left(\mu_{0,k}^{(N)} < w_{k,\varepsilon}^+\right) < \infty \quad (11)$$

then P -a.s. $\mathcal{N}_0(W) < \infty$.

Proof. By the strong law of large numbers, there is some $k_0 = k_0(\omega, \varepsilon)$ so that if $k \geq k_0$ and $x \in I_k$, then $(1 - \varepsilon)\alpha k \leq x$ in which case $W(x) \leq W((1 - \varepsilon)\alpha k)$. For such k , the Sturm oscillation and comparison theorems imply that $\mathcal{N}_{0,k,N}(W) \leq \mathcal{N}_{0,k,N}(W_\varepsilon^+ \mathbb{I}_k)$, i.e., the number of negative eigenvalues for the operator $H_{N,k} - W$ is not greater than the number of negative eigenvalues of $H_{N,k} - W_\varepsilon^+ \mathbb{I}_k$. The lowest eigenvalue for $H_{N,k} - W_\varepsilon^+ \mathbb{I}_k$ is $\mu_{0,k}^{(N)} - w_{k,\varepsilon}^+$ since $W_\varepsilon^+(x) = w_{k,\varepsilon}^+$ is constant for all $x \in I_k$. If (11) holds, then the Borel-Cantelli lemma implies that $\mu_{0,k}^{(N)} - w_{k,\varepsilon}^+ \geq 0$ P -a.s. for $k \geq k_1(\omega, \varepsilon) \geq k_0$. It means $\mathcal{N}_{0,k,N}(W_\varepsilon^+ \mathbb{I}_k) = 0$ and consequently $\mathcal{N}_{0,k,N}(W) = 0$ for all $k \geq k_1$. Dirichlet-Neumann bracketing implies $\mathcal{N}_0(W) \leq \mathcal{N}_{0,N}(W) = \sum \mathcal{N}_{0,k,N}(W) < \infty$. $\square$

Independence of the events $\{\mu_{0,k}^{(D)} < w_{k,\varepsilon}^-\}$ and similar ideas lead to the following theorem:

Theorem 4.2. *Let $\varepsilon > 0$ be given. If*

$$\sum_{k=1}^\infty P\left(\mu_{0,k}^{(D)} < w_{k,\varepsilon}^-\right) = \infty$$

then P -a.s. $\mathcal{N}_0(W) = \infty$.

In connection with the above theorems, we have the following lemmas:

Lemma 4.3. *For any $\varepsilon > 0$ let $w_{k,\varepsilon}^+ = W((1 - \varepsilon)\alpha k)$. There exists a positive constant c_N depending only on l and h such that $\mathcal{N}_0(W) < \infty$ P -a.s. whenever*

$$\sum_{k=1}^{\infty} P\left(L_k > \pi/\sqrt{w_{k,\varepsilon}^+} - c_N\right) < \infty \quad (12)$$

Proof. Suppose that P -a.s $\mathcal{N}_0(W) = \infty$. Then, by Theorem (4.1)

$$\sum_k P(\mu_{0,k}^{(N)} < w_{k,\varepsilon}^+) = \infty.$$

Since $\mu_{0,k}^{(N)}$ are independent random variables the second Borel-Cantelli lemma implies that P -a.s, there is a collection of intervals I_k with $k \rightarrow \infty$ on which $\sqrt{\mu_{0,k}^{(N)}} < \sqrt{w_{k,\varepsilon}^+}$. Therefore, $\sqrt{\mu_{0,k}^{(N)}} \rightarrow 0$ and by (9) it follows that $L_k \rightarrow \infty$. In this case by the asymptotic formula (10)

$$\frac{\pi}{L_k} \left(1 - \frac{B_{0,N}}{2L_k}\right) < \sqrt{w_{k,\varepsilon}^+}$$

for all k sufficiently large. Recall that $B_{0,N}$ is a positive constant depending only on l and h . This inequality and the fact that $L_k \rightarrow \infty$ immediately give

$$L_k \sqrt{w_{k,\varepsilon}^+} \geq \frac{1}{2}$$

for $k \gg 1$. Consequently,

$$L_k > \frac{\pi}{\sqrt{w_{k,\varepsilon}^+}} - \frac{\pi B_{0,N}}{2\sqrt{w_{k,\varepsilon}^+} L_k} \geq \frac{\pi}{\sqrt{w_{k,\varepsilon}^+}} - c_N$$

for sufficiently large k where c_N is a positive constant depending only on l and h . In this case

$$P(\mu_{0,k}^{(N)} < w_{k,\varepsilon}^+) = P\left(L_k > \pi/\sqrt{w_{k,\varepsilon}^+} - c_N\right)$$

so that $\sum_k P\left(L_k > \pi/\sqrt{w_{k,\varepsilon}^+} - c_N\right) = \infty$. □

Similar steps lead to the proof of the following lemma:

Lemma 4.4. *For any $\varepsilon > 0$ let $w_{k,\varepsilon}^- = W((1 + \varepsilon)\alpha k)$. There exists a positive constant c_D depending only on l and h such that $\mathcal{N}_0(W) = \infty$*

P-a.s. whenever

$$\sum_{k=1}^{\infty} P\left(L_k > \pi/\sqrt{w_{k,\varepsilon}^-} - c_D\right) = \infty$$

Remark. It follows that Theorems 1.3 and 1.4 presented in the introduction are a consequence of Lemmas 4.1 and 4.2 above. In fact, if $P(L_k > x) \sim \exp(-\eta x^\alpha/\alpha)$, then for $c = c_D$ or $c = c_N$, $P\left(L_k > \pi/\sqrt{w_{k,\varepsilon}^-} - c\right) \sim CP\left(L_k > \pi/\sqrt{w_{k,\varepsilon}^-}\right)$ for some positive constant C . The borderline for convergence or divergence of the sums $\sum P\left(L_k > \pi/\sqrt{w_{k,\varepsilon}^-} - c\right)$ is then achieved by taking $W(x) \sim C_{\eta,\alpha}/\ln^{2/\alpha} x$ where $C_{\eta,\alpha} = (\eta\alpha^{-1})^{2/\alpha}\pi^2$. In a similar way, if $P(L_k > x) \sim x^{-\alpha}$, then $P\left(L_k > \pi/\sqrt{w_{k,\varepsilon}^-} - c\right) \sim CP\left(L_k > \pi/\sqrt{w_{k,\varepsilon}^-}\right)$ for some positive constant C and the borderline between convergence or divergence is achieved if $W(x) \sim x^{-2/\alpha}$.

5. EXAMPLES

Our goal in this section is the proof of the following theorem.

Theorem 5.1. *Let $\{L_k\}$ be a sequence of independent, identically distributed non-negative random variables having exponential tails, that is,*

$$P\{L_k > x\} \sim e^{-\eta x}$$

*for some $\eta > 0$. Let $c_0 = \eta^2\pi^2$. If $W(x) < c_0/\ln^2 x$ for all $x \gg 1$, then $\mathcal{N}_0(W) < \infty$ *P*-a.s. If $W(x) > c_0/\ln^2 x$ for $x \gg 1$ then $\mathcal{N}_0(W) = \infty$ *P*-a.s.*

Proof. Suppose $W(x) \leq (c_0(1 - \delta))/\ln^2 x$ for some $0 < \delta < 1$ and $x \gg 1$. Then

$$P\{L_k > \pi/\sqrt{w_{k,\varepsilon}^+} - c_N\} \sim \frac{c}{k^{1+\beta}}$$

where $c = c(\eta, \varepsilon, \delta)$ and $1 + \beta = (1 - \delta)^{-1}$. Consequently,

$$\sum_{k=1}^{\infty} P\{L_k > \pi/\sqrt{w_{k,\varepsilon}^+} - c_N\} < \infty$$

and therefore by Lemma 4.3, $\mathcal{N}_0(W) < \infty$ *P*-a.s. On the other hand, if $W(x) \geq (c_0(1 + \delta))/\ln^2 x$ for $x \gg 1$, then

$$P\{L_k > \pi/\sqrt{w_{k,\varepsilon}^-} - c_D\} \geq \frac{c}{k^{1-\beta}}$$

where $1 - \beta = (1 + \delta)^{-1}$ and therefore

$$\sum_{k=1}^{\infty} P\{L_k > \pi/\sqrt{w_{k,\varepsilon}^-} - c_D\} = \infty.$$

By Lemma 4.4, $\mathcal{N}_0(W) = \infty$ P -a.s. $\square$

Example 5.2. Consider the Bernoulli potential $V_\omega(x) = h \sum \epsilon_k \xi_k(x)$ described in the introduction. By the Borel-Cantelli lemma, it follows that there exists P -a.s a sequence x_{n_k} of positive integers for which $V(x, \omega) = 0$ on $[x_k + 1/2, x_{k+1} - 1/2)$ and $L_k = x_{k+1} - x_k - 1 \rightarrow \infty$. Since for any $m \in \mathbb{Z}^+$, $P(L_k \geq m) = q^m = e^{-m \ln(1/q)}$ it follows that L_k have exponential tails. Hence by Theorem 5.1, the borderline between finite and infinite negative eigenvalues is given by a decay in $W(x)$ of the order $c_1/\ln^2 x$ with $c_1 = \pi^2 \ln^2(1/q)$.

REFERENCES

- [1] S. Molchanov and B. Vainberg, “Non-Random Perturbations of the Anderson Hamiltonian,” Journal of Spectral Theory, vol. 1, no. 2, 2011 pp. 179-195
- [2] M. Reed and B. Simon, “Methods of Modern Mathematical Physics, IV. Analysis of Operators,” Academic Press, Inc., London, 1978.
- [3] B. Levitan and I. Sargsjan, “Introduction to Spectral Theory: Selfadjoint Ordinary Differential Operators”, Translations of Mathematical Monographs, vol. 39, Amer. Math. Soc., Providence, Rhode Island 1975.